

***Vegetation management for optimizing conifer production in the northeast:
 lessons learned from research and practice.***

Doug Pitt, Canadian Forest Service, 1219 Queen St. E., Sault Ste. Marie, ON. P6A 2E5
 705-541-5610; dpitt@NRCan.gc.ca

Selected References

Conifer production/long-term responses

- Glover, G.R., and B.R. Zutter. 1993. Loblolly pine and mixed hardwood stand dynamics for 27 years following chemical, mechanical, and manual site preparation. *Canadian Journal of Forest Research* **23**: 2126-2132.
- Hearnden, K.W., S.V. Millson, and W.C. Wilson. 1992. A report on the status of forest regeneration. Ministry of Natural Resources, Forest Audit Secretariat, 70 Foster Drive, Suite 400, Sault Ste. Marie, ON P6A 6V5. 117 pp.
- MacLean, D.A. and M.G. Morgan. 1983. Long-term growth and yield response of young fir to manual and chemical release from shrub competition. *Forestry Chronicle*. **59**(4): 177-183.
- Sutton, R.F. 1995. White spruce establishment: initial fertilization, weed control, and irrigation evaluated after three decades. *New Forests* **9**: 123-133.

Additional:

- Borders, B.E., and R.L. Bailey. 2001. Loblolly pine – pushing the limits of growth. *Southern Journal of Applied Forestry* **25**(2): 69-74.
- Daggett, D.H. 2003. Long-term effects of herbicide and precommercial thinning treatments on species composition, stand structure, and net present value in spruce-fir stands in Maine: The Austin Pond study. M.Sc. Thesis, University of Maine, Orono, Maine, USA. 136 pp.
- Erdle, T., and J. Pollard. 2002. Are plantations changing the tree species composition of New Brunswick's forest? *Forestry Chronicle* **78**(6): 812-821.
- Oliver, W.W. 1990. Spacing and shrub competition influence 20-year development of planted ponderosa pine. *Western Journal of Applied Forestry* **5**(3): 79-82.
- Thompson, D.G., and D.G. Pitt. 2003. A review of Canadian forest vegetation management research and practice. *Annals of Forest Science* **60**: 559-572.
- Wagner, R.G., M. Newton, E.C. Cole, J.H. Miller, and B.D. Shiver. 2004. The role of herbicides for enhancing forest productivity and conserving land for biodiversity in North America. *Wildlife Society Bulletin* **32**(4): 1028-1041.

Degree of conifer response is a function of treatment intensity

- Glover, G.R., and B.R. Zutter. 1993. Loblolly pine and mixed hardwood stand dynamics for 27 years following chemical, mechanical, and manual site preparation. *Canadian Journal of Forest Research* **23**: 2126-2132.
- Pitt, D.G., and F.W. Bell. 2005. Juvenile response to conifer release alternatives on aspen-white spruce boreal mixedwood sites. Part I: Stand structure and composition. *Forestry Chronicle* **81**(4): 538-547.
- Pitt, D.G., M. Mihajlovich, and L.M. Proudfoot. 2004. Juvenile stand responses and potential outcomes of conifer release efforts on Alberta's spruce-aspen mixedwood sites. *Forestry Chronicle* **80**(5):583-597.
- Pitt, D.G., R.G. Wagner, and W.D. Towill. 2004. Ten years of vegetation succession following ground-applied release treatments in young black spruce plantations. *Northern Journal of Applied Forestry* **21**(3):123-133.

Additional:

Greifenhagen, S., D.G. Pitt, M.C. Wester, and F.W. Bell. 2005. Juvenile response to conifer release alternatives on aspen-white spruce boreal mixedwood sites. Part II: Quality of aspen regeneration. *Forestry Chronicle* 81(4): 548-558.

Pitt, D.G., A.E. Morneau, P. Bunce, and F.W. Bell. Five years of vegetation succession following vegetation management treatments in a jack pine ecosystem. *Northern Journal of Applied Forestry* 17(3): 100-109.

Early action is critical

Pitt, D.G., C.S. Krishka, F.W. Bell, and A. Lehela. 1999. Five-year performance of three conifer stock types on fine sandy loam soils treated with hexazinone. *Northern Journal of Applied Forestry* 16(2): 72-81.

Pitt, D.G., B. Parker, A. Morneau, A. Stinson, W. Bell, and L. Lanteigne. 2004. Partner's Report, 2004 field season – Effects of early herbaceous and woody vegetation control on eastern white pine. Natural Resources Canada, Canadian Forest Service, 1219 Queen St. E., Sault Ste. Marie, ON. 30 p. (http://www.forestresearch.ca/product_catalogue/reports.htm, report RP-25).

Pitt, D.G., B. Parker, A. Morneau, A. Stinson, C. Yurich, W. Bell, and L. Lanteigne. 2005. Partner's Report, 2005 field season – Effects of early herbaceous and woody vegetation control on eastern white pine. Natural Resources Canada, Canadian Forest Service, 1219 Queen St. E., Sault Ste. Marie, ON. 30 p. (http://www.forestresearch.ca/product_catalogue/reports.htm, report RP-31).

Wagner, R.G., G. H. Mohammed, and T.L. Noland. 1999. Critical period of interspecific competition for northern conifers associated with herbaceous vegetation. *Canadian Journal of Forest Research* 29: 890-897.

Wagner, R.G. and A.P. Robinson. 2006. Critical period of interspecific competition for four northern conifers: Ten-year growth response and associated vegetation dynamics. *Canadian Journal of Forest Research* (in press).

Wood, J.E., and F.W. von Althen. 1993. Establishment of white spruce and black spruce in boreal Ontario: Effects of chemical site preparation and post-planting weed control. *Forestry Chronicle* 69(5): 554-560.

Additional:

Shiver, B.D., and S.W. Martin. 2002. Twelve-year results of a loblolly pine site preparation study in the Piedmont and Upper Coastal Plain of South Carolina, Georgia, and Alabama. *Southern Journal of Forest Research* 26(1): 32-36.

Herbaceous weed control is important

Jaciw, P. 1972. How silvicides gave 10-year assist to conifer regeneration. *Can. For. Ind.* 92: 30-36

Miller, J.H., B.R. Zutter, S.M. Zedaker, M.B. Edwards, and R.A. Newbold. 2003. Growth and yield relative to competition for loblolly pine plantations to midrotation – a southeastern United States regional study. *Southern Journal of Applied Forestry*. 27(4): 237-252.

Pitt, D.G., B. Parker, A. Morneau, A. Stinson, W. Bell, and L. Lanteigne. 2004. Partner's Report, 2004 field season – Effects of early herbaceous and woody vegetation control on eastern white pine. Natural Resources Canada, Canadian Forest Service, 1219 Queen St. E., Sault Ste. Marie, ON. 30 p. (http://www.forestresearch.ca/product_catalogue/reports.htm, report RP-25).

Pitt, D.G., B. Parker, A. Morneau, A. Stinson, C. Yurich, W. Bell, and L. Lanteigne. 2005. Partner's Report, 2005 field season – Effects of early herbaceous and woody vegetation control on eastern white pine. Natural Resources Canada, Canadian Forest Service, 1219 Queen St. E., Sault Ste. Marie, ON. 30 p. (http://www.forestresearch.ca/product_catalogue/reports.htm, report RP-31).

Pitt, D.G., R.G. Wagner, and W.D. Towill. 2004. Ten years of vegetation succession following ground-applied release treatments in young black spruce plantations. *Northern Journal of Applied Forestry* 21(3):123-133.

Additional:

- Glover, G.R., J.L. Creighton, and D.H. Gjerstad. 1989. Herbaceous weed control increases loblolly pine growth: A 12-year study on three locations demonstrated consistent and substantial volume gain. *Journal of Forestry* 87(2): 47-50.
- Harrington, T.B., R.G. Wagner, S.R. Radosevich, and J. D. Walstad. 1995. Interspecific competition and herbicide injury influence 10-year responses of coastal Douglas-fir and associated vegetation to release treatments. *Forest Ecology and Management* 76: 55-67.
- Haywood, J.D. 1994. Tenth-year results of herbaceous weed control in a loblolly pine plantation. *Southern Journal of Applied Forestry* 18(3): 105-109.
- Haywood, J.D., and A.E. Tiarks. 1990. Eleventh-year results of fertilization, herbaceous, and woody plant control in a loblolly pine plantation. *Southern Journal of Applied Forestry* 14(4): 173-177.
- Lauer, D.K., G.R. Glover, and D.H. Gjerstad. 1993. Comparison of duration and method of herbaceous weed control on loblolly pine response through mid rotation. *Canadian Journal of Forest Research* 23: 2116-2125.
- Monleon, V.J., M. Newton, C. Hooper, and J.C. Tappeiner. 1999. Ten-year growth response of young Douglas-fir to variable density varnishleaf ceanothus and herb competition. *Western Journal of Applied Forestry* 14(4): 208-213.
- Nelson, L.R., B.R. Zutter, and D.H. Gjerstad. 1985. Planted longleaf pine seedlings respond to herbaceous weed control using herbicides. *Southern Journal of Applied Forestry* 9(4): 236-240.
- Pitt, D.G., C.S. Krishka, F.W. Bell, and A. Lehela. 1999. Five-year performance of three conifer stock types on fine sandy loams treated with hexazinone. *Northern Journal of Applied Forestry* 16(2): 72-81.
- Zutter, B.R., G.R. Glover, and D.H. Gjerstad. 1987. Vegetation response to intensity of herbaceous weed control in a newly planted loblolly pine plantation. *New Forests* 4: 257-271.
- Zutter, B.R., and J.H. Miller. 1998. Eleventh-year response of loblolly pine and competing vegetation to woody and herbaceous plant control on a Georgia flatwoods site. *Southern Journal of Applied Forestry* 22: 88-95.